

Leaflet FRUITER



COMPOSITION:

Potassium Oxide (K₂O): 18% w/w

Boron: 0.2% w/w

Manitol: 0.1%

DOSING INSTRUCTIONS:

Crop	Application stage	Dosage in L / ha
Vegetables	Every 10-12 days from start of fruit development	2 - 3
Fruit trees	2-4 applications from fruit set to ripening	2 - 3
Flowers	During the flowering stage	2 - 3
Cereals	During the grain filling stage	1 - 1.5
Arable crops	During the grain filling stage	1 - 1.5
Tropical fruits	From the start of fruit development and ripening stage	2 - 3

The indicated dosages and application stages are subject to soil and climat conditions, influence of previous crops and other specific conditions. Exact dosages and application stages can only be given after an objective diagnostic procedure.

It is advised to conduct a compatibility test before mixing with other products.

PRODUCT CHARACTERISTICS:

Naturally sourced raw material to create the perfect bio-stimulant product. Increases crop resilience, improves plant health and boost its natural processes.

FRUITER promotes the increasing of sugar content in the fruit, thus improving its quality and appearance. It also improves fruit firmness, increases the fruit size and earliness.

The resulting products will have:

- Better taste
- More appealing appearance
- Greater shelf life
- Earlier harvest

CAUTION AND STORAGE:

Shake well before each use. Keep out of reach of children. Avoid contact with eyes, skin and clothing. Avoid breathing spray mist. If in eyes, flush immediately with running water for at least 15 minutes keeping eyelids open. If on skin, wash thoroughly with running water and non-abrasive soap. If ingested, immediately drink several glasses of water or milk. If irritation persists, seek medical attention.

Store this product in original package, in area of low fire risk, away from heat, sources of ignition and combustible materials. Keep lid tightly closed.