

Leaflet ROOTER



COMPOSITION:

Total Nitrogen (N): 2.7% w/w

Organic Nitrogen: 2.4% w/w

Ammoniacal Nitrogen: 0.3% w/w

Total Potassium (K₂O): 3.3% w/w

Organic Carbon: 17.4% w/w

DOSING INSTRUCTIONS:

Crop	Application stage	Dosage in L / ha
Vegetables	Every 7-10 days, starting 2-3 weeks after planting	3 - 5
Fruit trees	Every 15-25 days from the beginning of the season	7 - 10
Flowers	Every 7-10 days, starting 2-3 weeks after planting	3 - 5
Cereals	With first herbicide application	1 - 2
Arable crops	With first herbicide application	1 - 2
Tropical fruits	Every 15-25 days from the beginning of the season	7 - 10

The indicated dosages and application stages are subject to soil and climat conditions, influence of previous crops and other specific conditions. Exact dosages and application stages can only be given after an objective diagnostic procedure.

It is advised to conduct a compatibility test before mixing with other products.

PRODUCT CHARACTERISTICS:

Naturally sourced raw material to create the perfect bio-stimulant product. Increases crop resilience, improves plant health and boost its natural processes.

ROOTER promotes root development and aids the plant to better tolerate abiotic stress. As a result, the plants will:

- Quickly develop a larger root system
- Increase their nutrient absorption capacity
- Increase their drought tolerance
- Enhance fruit uniformity

CAUTION AND STORAGE:

Shake well before each use. Keep out of reach of children. Avoid contact with eyes, skin and clothing. Avoid breathing spray mist. If in eyes, flush immediately with running water for at least 15 minutes keeping eyelids open. If on skin, wash thoroughly with running water and non-abrasive soap. If ingested, immediately drink several glasses of water or milk. If irritation persists, seek medical attention.

Store this product in original package, in area of low fire risk, away from heat, sources of ignition and combustible materials. Keep lid tightly closed.